

2 GRAM SODIUM DIET

WHAT YOU SHOULD KNOW:

A 2 gram sodium diet limits high sodium foods in your diet. No table salt is allowed at meals or during cooking while you are on this diet. The amount of milk is also limited because of the amount of sodium it contains. A high amount of sodium in your diet can make your blood pressure go up and can cause other health problems. The goal of a 2 gram sodium diet is to prevent or lower high blood pressure. This diet can also keep your body from holding extra fluid. Problems with your liver and kidneys are another reason to follow this diet.

INSTRUCTIONS:

How can I use food labels to choose foods that are low in sodium? Reading food labels is a good way to learn how much sodium is in foods. Ask your caregiver for more information about how to read food labels. Food labels list the amount of sodium in the food in milligrams. **Avoid foods that contain more than 500 mg of sodium in one serving.** Buy low-sodium substitutes for the foods you enjoy. Following are some words about sodium that may appear on a label.

- **Sodium-free:** Less than five mg in each serving.
- **Very low sodium:** 35 mg of sodium or less in each serving.
- **Low sodium:** 140 mg of sodium or less in each serving.
- **Reduced sodium:** At least 25 percent less sodium in each serving. For example, if the food usually has 800 mg of sodium, the same food made with reduced sodium would contain 600 mg of sodium.
- **Light in sodium:** Fifty percent less sodium in each serving. For example, if the food usually has 500 mg of sodium in each serving, the same food prepared "light in sodium" would have 250 mg of sodium.
- **Unsalted, No added salt, and Without added salt:** No salt is added during processing.
- **Lightly salted:** Fifty percent less sodium has been added to the food than would normally be added. For example, if 1000 mg of sodium were normally added, only 500 mg would be added to a food that is "lightly salted".

What should I avoid eating and drinking while on a 2 gram sodium diet? **Breads, cereals, rice and pasta:**

- Breads, rolls and crackers with salted tops.
- Quick breads.
- Instant hot cereals.
- Pancakes.

- Commercial (store-bought) bread stuffing, self-rising flour and biscuit mixes.
- Commercial breadcrumbs, cracker crumbs, or bread stuffing.
- Rice or pasta mixes.

Fruits and vegetables:

- Fruits processed with salt or sodium-containing ingredients, such as some dried fruits.
- Regular canned vegetables, and olives.
- Sauerkraut, pickled vegetables, and others prepared in brine.
- Frozen vegetables in sauces.
- Vegetables seasoned with ham, bacon, or salt pork.
- Commercially prepared potato mixes.

Meats and meat products:

- Smoked, cured, salted, koshered, or canned meat, fish or poultry, including bacon, chipped beef, cold cuts, ham, hot dogs, sausage, sardines, anchovies, crab, lobster, imitation seafood, marinated herring, and pickled meats and frozen breaded meats.
- Pickled eggs.
- Processed cheese, cheese spreads, and sauces.
- Salted nuts.

Beverages and soups:

- Malted milk, milkshake and chocolate milk.
- Regular vegetable or tomato juice.
- Commercially softened water (for drinking or cooking).
- Regular canned or dehydrated soups.
- Broths and bouillon.

Fats, desserts and sweets:

- Regular salad dressings containing bacon fat, bacon bits, and salt pork.
- Snack dips made with instant soup mixes or processed cheese.

- Instant pudding mixes and cake mixes.

Seasonings, sauces, mixes, and other foods:

- Seasoning made with salt including garlic salt, celery salt, onion salt, and seasoned salt.
- Sea salt, rock salt, kosher salt, meat tenderizers and monosodium glutamate (MSG).
- Regular soy sauce, barbecue sauce, teriyaki sauce, steak sauce, Worcestershire sauce, and most flavored vinegars.
- Canned gravy and mixes, regular condiments and salted snack foods.

What other guidelines should I follow while on a 2 gram sodium diet?

- Avoid using salt and ingredients such as baking soda and soy sauce that are high in sodium. Be careful to avoid these during food preparation, and at the table. Meals eaten at restaurants, especially fast food restaurants are often high in sodium. Some restaurants have nutrition information that tells you the amount of sodium in their foods. When ordering food at a restaurant, ask your waiter to prepare your food with less, or no salt.
- Talk with your caregiver about using salt substitutes. Some salt substitutes have ingredients that may change the way some of your medicines work. Check with your caregiver or pharmacist about products that may contain sodium. Some of these products include antacids, medicine, toothpaste and chewing tobacco.

Risks

- Eating more than two grams of sodium in a day may cause your body to hold extra fluid, causing you to gain weight. It may make your blood pressure higher. Eating more than two grams of sodium in a day may change the way your medicines work.
- Very rarely, a two gram sodium diet may cause the amount of sodium in your blood to be too low. Low amounts of sodium in your blood may cause nausea, confusion, and make you less alert. **Call your caregiver right away if you have any of these symptoms.**

© 1974-2007 Thomson MICROMEDEX. All rights reserved.

**2 GRAM SODIUM DIET - AfterCare(R) Instructions(ER/ED), English
Thursday, January 4, 2007 9:00:47 AM**

*THIS INFORMATION IS MADE AVAILABLE FOR YOUR USE BY
ST. JOHN'S HOSPITAL, HEALTH SCIENCES LIBRARY
800 E Carpenter, Springfield, IL 62769 (217-544-6464).
Visit our Web site at <http://www.st-johns.org>.*

DISCLAIMER: This information is for informational purposes only and is not designed to take the place of medical or other professional advice. If professional advice is required, the services of a physician or other medical professional should be sought. By accessing this information, the user acknowledges that St. John's Hospital and its affiliates shall not be liable for any damage, claim, or loss that may arise from the use of this information or for any diagnosis, treatment, or other action taken in reliance